

## Michigan TRAUMA AND ENVIRONMENTAL **BLEEDING CONTROL (BCON)**

Initial Date: 3/23/2018

Revised Date: -08/23/2024<del>05/23/2023</del> Section: 2-13

## Bleeding Control

## Indications:

Patients with significant traumatic or non-traumatic (i.e., hemodialysis access) external hemorrhage

- 1. Follow General Pre-hospital Care-Treatment Protocol and Soft Tissue & **Orthopedic Injuries-Treatment Protocol.**
- Apply direct pressure to the wound with clean gauze using universal precautions. 2.
- If the bleeding is not controlled with direct pressure, treat according to the 3. location of the wound.
  - a. Extremity bleeding apply tourniquet:(Refer to Tourniquet Application-**Procedure Protocol**)
    - i. If tourniquet unsuccessful apply second/adjacent tourniquet per Tourniquet Application-Procedure Protocol.
    - ii. NOTE- tourniquet may be painful, see Pain Management-Procedure Protocol.
  - b. <u>Junctional hemorrhage (Nneck, axilla/shoulder or groin bleeding)</u>:
    - i. Pack wound with MCA approved hemostatic dressing (if available and, following manufacturer's instructions) or clean gauze.
    - ii. Use as much of the dressing/gauze as needed to stop the blood flow.
    - iii. Quickly apply pressure until the bleeding stops. (Approximately 3-5 minutes)
    - iv. Leave the dressing in place and wrap area with bandaging to secure the dressing.
    - v. Caution must be exercised when packing neck wounds to avoid airway compromise. Avoid an encircling bandage of the neck.
    - iv.vi. Packing penetrating head, chest, and abdominal wounds is contraindicated.
- Do not remove the bandage or hemostatic dressing/gauze 4.
- 5. Elevate the injury, if possible.
- Reassess for bleeding through or around the dressing. 6.
- For patients who have signs or symptoms of shock, secondary to hemorrhage, 7. refer to Hemorrhagic Shock-Treatment Protocol.
- Transport according to Adult and Pediatric Trauma Triage-Treatment Protocol 8. and MCA Transport Protocol



## Notes:

If hemostatic dressing is used, contact medical control to advise of application, document time of use, and send packaging from dressing to hospital with patient for removal instructions.