

Michigan
COMMUNITY INTEGRATED PARAMEDICINE
Treatment Protocol
DIABETIC CARE

Initial Date: November 19, 2020

Revised Date:

Section 11-51

This protocol is for trained CIP Paramedics only. If during assessment, procedure, or treatment the patient is found to have a medical emergency in need of hospital treatment, the CIP visit will be suspended, and local MCA protocols utilized.

Purpose: To provide additional guidelines beyond the CIP General Assessment and Care for evaluating, caring for and educating patients with Diabetes.

- I. Follow **CIP Patient General Assessment and Care protocol**
- II. Obtain additional vital signs, history, and diagnostics pertinent to condition and/or as ordered by physician.
 - a. Vitals: BGL
 - b. History: Last oral intake, diet, medication changes and compliance
- III. On-scene medication administration may include:
 - a. Use of approved MCA protocols and medications up to the extent of standard paramedic treatment according to protocol.
 - b. Assist with patient's prescribed home medications that are not included in standard EMS treatment protocols
 - c. Use of patient's prescribed medications beyond the parameters of the prescription must have direct consultation with the referring physician prior to administration
 - d. Oral high caloric fluid
 - e. Oral glucose gel or tablets
 - f. IV Fluid bolus of 0.9% NS maximum dose 2L
- IV. On-scene education and suggested support sources may include:
 - i. Diabetes Self-Management Education

MCA Name: [Click here to enter text.](#)

MCA Board Approval Date: [Click here to enter text.](#)

MCA Implementation Date: [Click here to enter text.](#)

Protocol Source/References: [Click here to enter text.](#)